# **Elephant Necropsy Protocol**

(Elephas maximus and Loxodonta africana)

The American Zoo and Aquarium Association
Elephant Species Survival Plan
and
The Elephant Research Foundation

Revised 1997

# **Table of Contents**

Introduction	1
Equipment Checklist	2
Logistics / Necropsy Tips	2
Gross Examination Worksheet	3-5
Measurements	6-10
Search List	10-1
Selected Bibliography	11-12
Researchers Interested in Participating in Necropsies	13
Research Requests.	13-19

#### Introduction

This protocol is a cooperative effort of the Elephant Species Survival Plan (SSP) Propagation Group of the American Zoo and Aquarium Association (AZA) and the Elephant Research Foundation (ERF). Its purpose is to provide a format for the systematic collection of information and samples that will add to our knowledge of elephants. All North American institutions holding elephants will have received a copy.

We hope that most institutions will not have to face the immense task of performing an elephant necropsy, but should a death occur, it should be viewed as an important learning opportunity. Although it may not be feasible to collect all the information and samples requested, we encourage the collection of as much as possible.

The Search List describes those parts of the anatomy for which data is lacking or about which previous observations need to be confirmed or refuted. The Measurements Checklist may seem tedious, but only this type of attention to detail will allow us to expand our knowledge of elephant anatomy. Both of these requested data sets are optional.

Acquainting oneself with the protocol and having the necessary equipment ready will facilitate sample collection. It is suggested that a necropsy team be designated in advance; the ability to mobilize skilled individuals quickly will save valuable time particularly in the event of a sudden death. Veterinarians, anatomists, and pathologists from nearby universities may be enlisted to assist the institution's staff. In addition, a list of researchers interested in participating in elephant necropsies is included in this protocol.

A revised Research Request List will be forwarded periodically as new requests are received and projects end. Contact Susan Mikota for current requests. A copy of the completed protocol together with the histopathology and any other lab reports should be forwarded to Dr. Mikota.

Susan Mikota D.V.M., Director of Veterinary Research and Animal Health Audubon Center for Research of Endangered Species & Freeport-McMoran Audubon Species Survival Center 14001 River Road New Orleans, LA 70131

Work: (504) 398-3111 Fax: (504) 398-3100

email: smikota@acres.org

Richard Montali D.V.M., Head Department of Pathology National Zoo 3000 Block of Connecticut Ave. NW Washington DC 20008-2598

Work: (202) 673-4869 Fax: (202) 673-4660

email: nzpem026@sivm.si.edu

### **Equipment Checklist**

- 1. Standard large animal necropsy instruments. Multiple scalpel handles, duplicates or triplicates of other instruments. Extra box of scalpel blades, knife sharpener.
- 2. Retractors of various sizes and shapes. Self-retaining retractors with one or two movable arms mounted on a slide bar are most useful.
- 3. Sterile instruments for culture collection.
- 4. 10% neutral buffered formalin.
- 5. 10% glutaraldehyde (for electron microscopy).
- 6. Containers for sample collection.
- 7. Culture swabs, sterile urine cups, glass slides.
- 8. Serum tubes for blood and urine collection.
- 9. Aluminum foil and plastic bags for freezing tissues.
- 10. Labels and waterproof marking pens.
- 11. Scale for obtaining organ weights.
- 12. Tape measure (metric), at least 2 meters long.
- 13. Chain saw and axe to cut through the cranium. Hammers, chisels and hand saws.
- 14. Hoist/crane.
- 15. Carts on rollers to move heavy parts.
- 16. Coveralls, boots, gloves, caps, masks, protective eye and head gear.
- 17. Accessible water supply with hose.
- 18. Camera and film, extra batteries.
- 19. First aid kit.

### Logistics and Necropsy Tips

Heavy equipment may be necessary to move a dead elephant. For an on site necropsy, chains and a tow truck may be sufficient to reposition the animal or to move it a short distance. If the animal must be transported to a remote site, a truck with a hoist will be needed. It may be easier to manipulate the animal onto a flatbed trailer. Vehicles must be able to handle these approximate weights: female Asian: 2,300 - 3,700 kg; male Asian: 3,700 - 4,500 kg; female African: 2,300 - 4,000 kg; male African: 4,100 - 5,000 kg. Trucks can generally be rented from equipment rental companies, or may be available from the telephone company. If a flatbed carrier is used the animal will need to be strapped to the bed and covered with a tarp (a baseball diamond infield tarp works well). If transportation will be delayed, the carcass can be covered with ice.

If death is imminent or euthanasia is planned, completion of the measurement checklist antemortem will save time at necropsy. Otherwise, measurements should be done as soon after death as possible.

Assigning specific tasks to team members will help the necropsy to proceed in an orderly manner. For example, a team may be assigned to each of these areas: head, forelegs, hindlegs, abdominal region. One person should oversee the collection, labeling, and processing of research materials and any communication concerning research request. It may be helpful to designate a media spokesperson.

Dissection of the head is best completed after separating it from the body. A good portion of the cranium must be damaged to remove the brain intact; a chain saw, large axe, and chisels are needed to penetrate the thick cranium. A posterior approach to brain removal can be made by 3 connecting deep cuts with a chain saw in the margins of the flattened triangle formed at the base of the elephant skull. Then remove the bony plate in chunks with a curved crow-bar. Use of a chain saw on bone can be hazardous and cause shrapnel-like fragments to be launched. Protective head and face gear should be worn by the chain saw operator and attending personnel removed from the immediate area.

Dissection of the thoracic cavity is best performed by at least two people. After the initial incision at the ventral midline is made, one person holds the retractor and the other cuts the tensed skin. Once the sternum is exposed, the ribs are separated at the cartilaginous attachment and adjustable retractors are applied to hold the cavity open. The heart, lungs, and associated structures may be removed "en bloc" with the diaphragm. Visceral and parietal pleura are normally adhered; there is no pleural space.

# SSP Elephant Necropsy Protocol Gross Examination Worksheet

Institution/Owner				<u> </u>	
Address		Country			
Species	Id#	Isis#	Studbook#	<u> </u>	
Birth Date/Age	Se	xWeig	ht (KG)	(Actual/Estimate)	
Death Date	Death Lo	ocation			
Necropsy Date	Necropsy Location		PM Into	erval	
Captive Born □	Wild Caught [	]	<del></del>		
History (Include clinica	al signs, circumstances of c	leath, clinical lab w	ork, diet & housing)		
, •	GROSS s are noted mark as norn al and nutritional condition,		ed (NE); use addition		
Musculoskeletal Syst	em (Bones, marrow, joints,	muscles)			
Body Cavities (Fat sto	res, pleura, thymus, lymph n	odes)			
Spleen					
Respiratory System (1	Nasal passages, pharynx, lary	vnx, trachea, bronchi,	lungs, regional lymph	nodes)	

Cardiovascular System (heart, pericardial sac, great vessels, myocardium, valves, chambers)	
Digestive System (mouth, teeth, tongue, esophagus, stomach, small intestine, cecum, large intestine, gallbladder, pancreas, mesenteric lymph nodes)	anus, liver and
Urinary System (kidneys, ureters, bladder, urethra)	
Reproductive System (testes/ovaries, uterus & cervix, penis/vagina, urogenital canal, accessory sex orgland, placenta)	gans, mammary
Endocrine System (thyroids, parathyroids, adrenals, pituitary)	
Central Nervous System (brain, meninges, spinal cord)	
Sensory Organs (eyes, ears)	<u>:</u>
Additional Comments or Observations:	
Prosector:Date:	
Summarize Preliminary Diagnoses:	
Laboratory Studies: Results of cytology, fluid analysis, urinalysis, serum chemistries, bacteriology, parasitology, x-ray, photography, other	ogy, mycology,


## Measurements (Optional)

This data sheet is a general guideline to the pre-euthanasia or post mortem measuring of an elephant. Refer to the anatomical diagrams, Figures 1 and 2. The numbering system begins at the trunk and continues in a clockwise direction. All measurements should be taken in a straight line, except when indicated otherwise. Measurements to be taken between corresponding points on opposite sides of the body are marked with a plus symbol (+). These should be taken in a straight line, essentially through, not around, the elephant. Calipers can be improvised from two long straight poles or straight edges. Place the end of each pole on one of the two points, keeping the poles parallel to one another. Measure the straight line distance between the free ends of the two parallel poles.

	GENERAL		
Subject		Reference numbers on figures	Measurement between these points (cm)
Tip of trunk to	tip of tail (along the curve)	Fig. 1: 1-9	
Length of trun	k (along the curve)	Fig. 1: 1-2	
Length of tail		Fig. 1: 8-9	
Shoulder heig	ht	Fig. 1: 5-14	
Dorsum heigh	t (the highest point of back or "hump")	Fig. 1: 6-13	
	DETAILED		
TRUNK	Tip to base length	Fig. 1: 1-2	
	Tip width	Fig. 1: 1-21	
	Base width	Fig. 1: 40+	
HEAD	Dorsal length (along the curve)	Fig. 1: 2-3	
	Ventral length (along the curve)	Fig. 1: 19-20	
	Neck height	Fig. 1: 3-19	
	Width between ears	Fig. 1: 38+	
	Width between temporal glands	Fig. 1: 38a+	
	Width between eyes	Fig. 1: 39+	
	Width of mouth	Fig. 1: 40+	
EAR	Anterior width	Fig. 1: 22-23	
	Posterior width	Fig. 1: 24-25	
	Dorsal length	Fig. 1: 22-24	
	Ventral length	Fig. 1: 23-25	

	Subject	Reference numbers on figures	Measurement between these points (cm)
NECK	Length	Fig. 1: 3-4	
	Width	Fig. 1: 37+	
	Height	Fig. 1: 3-19	
BODY	Dorsal length (along the curve; number 7 is in a straight line with number 11)	Fig. 1: 4-7	
	Middle length (make sure this and the next measurement are taken parallel to each other)	Fig. 1: 32-26	
	Bottom length (make sure this and the previous measurement are taken parallel to each other)	Fig. 1: 10-27-18a	
	Width at front	Fig. 1: 36+	
	Width at middle	Fig. 1: 35+	
	Width at back	Fig. 1: 34+	
	Height at front of forelimb	Fig. 1: 5-27	
	Height at front of hindlimb	Fig. 1: 6-30	
	Height at back of hindlimb	Fig. 1: 7-10	
TAIL	Length (excluding hairs)	Fig. 1: 8-9	
	Width at base	Fig. 1: 8-33	, the second second
FORELIMB	Length (height)	Fig. 1: 16-26	
	Width at top	Fig. 1: 18b-28	
	Width at bottom (include side width, if different)	Fig. 1: 15-17	
HINDLIMB	Length (height)	Fig. 1: 11-31	
	Width at top	Fig. 1: 29-31	
	Width at bottom (include side width, since it is narrower)	Fig. 1: 11-12	

	Subject	Reference numbers on figures	Measurement between these points (cm)
FEET	Count number of "toenails"		Left front
			Right front
			Left hind
			Right hind
TEETH	Total number of plates (including very small ones)	Figure 2a: (see 1b)	
(can be	Total length	Fig. 2a: 1	
measured	Maximum width	Fig. 2a: 3 (see 1a)	
soon after death or at a later date)	Maximum grinding length of individual teeth	Fig. 2a: 1a	
	Maximum grinding length of entire grinding surface	Fig. 2a: 1b	
	Maximum height	Fig. 2a: 2	
	Weight (in grams)		
TUSKS	Present Absent	Figure 2b	
	Length from tip to gum line	Fig. 2b: b-c	
,	Length from gum line to base	Fig. 2b: a-c	
	Length of pulp cavity	Fig. 2b: a-d	
	Width of pulp cavity	Fig. 2b: e-e	· Š
	Total length	Fig. 2b: a-b	
PENIS	Circumference at base		
	Circumference at head		
	Length		
CLITORIS	Circumference at base		
	Circumference at head		






Figure 1. Generalized illustrations of an elephant showing points for measurement; a) after Deraniyagala (1955); all others by Shoshani. Letter "V" on the head indicates the approximate location of the vent gland.




Figure 2. Generalized illustrations of the teeth and tusk showing points for measurement. After Roth and Shoshani (1988).

### Search List (Optional)

The following are anatomical features that need to be confirmed or refuted, or for which few data exist. They are not arranged in order of importance, but rather as one studies the elephant by regions from the tip of the trunk to the tip of the tail. Please be aware of these anatomical questions and attempt to obtain the needed additional data as you proceed in your dissection.

- 1. Record the number of toes.
- 2. Weigh skin after dissection from limbs and carcass. Optional.
- 3. Search for sesamoids especially under tendons. There may be one at the proximal end of the humerus, but check other sites as well.
- 4. Obtain total skeletal weight. Remove as much soft tissue as possible.
- Note any pathological conditions in the joints. Slight erosions on articular surfaces can be viewed best in fresh tissues and should be examined soon after death. Grooves and fractures on articular surfaces cannot be mistaken and should be sought. Look also for "joint mice," calcium deposits, and any other abnormal signs.

- 6. Measure the **volume of the nasal passages** by instilling water soon after death or by measuring the diameter of the passages at intervals (record total length of trunk and diameter of passages at intervals of 10 cm).
- 7. Look for the intercommunicating canal between the two nasal passages of the trunk and the associated fibrous arches by sectioning the trunk every 10-20 cm. These structures were described as being located 13 cm from the tip of the trunk in a young female Asian elephant. Other searches in adult Asian females have revealed neither the arches nor the canals (Shoshani et al., 1982).
- 8. Harvest the lenses from the eyes and weigh them (or keep intact eyes frozen).
- 9. Search for the **trachea-esophageal muscle**. This muscle is small and may be overlooked or cut during dissection so we suggest that a section about 20 cm posterior and 50 cm or more anterior to the bifurcation be removed and examined carefully outside the carcass. This muscle was found in only three of twelve elephants examined (Shoshani *et al.*, 1982).
- 10. Examine the **dividing arrangement of the arteries from the aortic arch**. There are two possibilities: three branches or two branches. In the three-branch arrangement the sequence is right subclavian, a trunk common to the two carotids and the left subclavian. In the two-branch arrangement, the right subclavian and the common carotids merge into one vessel and the left subclavian remains separate.

## Literature Cited and Selected Bibliography

- Benedict, F.G. (1936). The physiology of the elephant. Carnegie Institution of Washington.
- Blair, P. (1710a). Osteographia elephantina: or, a full and exact description of all the bones of an elephant which dy'd near Dundee, April the 27th, 1706, with their several dimensions, etc. Part I. *Philos Trans R Soc Lond [Biol]* 27(326):51-116.
- Blair, P. (1710b). Osteographia elephantina: or, a full and exact description of all the bones of an elephant which dy'd near Dundee, April the 27th, 1706, with their several dimensions, etc. Part II. *Philos Trans R Soc Lond [Biol]* 27(327):117-168.
- Deraniyagala, P.E.P. (1955). Some extinct elephants, their relatives and the two living species. Ceylon National Museum Publ., Colombo.
- Eales, N.B. (1925). External features, skin, and temporal gland of a foetal African elephant. *Proc Zool Soc Lond* 2:445-456.
- Eales, N.B. (1926). The anatomy of the head of a foetal African elephant, *Elephas africanus* (Loxodonta africana). Trans Roy Soc Edin 54, part III (11):491-551 plus 12 plates.
- Eales, N.B. (1928). The anatomy of a foetal African elephant, *Elephas africanus* (*Loxodonta africana*) Part II. The body muscles. *Trans Roy Soc Edin* 55, part III (25):609-642 plus 5 plates.
- Eales, N. (1929). The anatomy of a foetal African elephant, *Elephas africanus* (*Loxodonta africana*). Part III. The contents of the thorax and abdomen, and the skeleton. *Trans Roy Soc Edin* 56, part I (11):203-246.
- Hill, W.C.O. (1938). Studies on the cardiac anatomy of the elephant: II -- the heart and great vessels of a foetal Asiatic elephant. *J Sci* 21:44-61.
- Laursen, L. and Beckoff, M. (1978). Loxodonta africana. Mamm Species 92:1-8.
- Laws, R.M. (1966). Age criteria for the African elephant, Loxodonta a. africana. East Afr Wildl J 4:1-37.
- Laws, R.M. (1967). Eye lens weight and age in the African elephant. East Afr Wildl J 5:46-52.
- Laws, R.M., Parker, I.S.C. and Archer, A.L. (1967). Estimating live weights of elephants from hindleg weights. East Afr Wildl J 5:106-111.
- Mariappa, D. (1986). Anatomy and histology of the Indian elephant. Indira Publishing House, Oak Park, MI.
- Miall, L.C. and Greenwood, F. (1879a). The anatomy of the Indian elephant. Part I. The muscles of the extremities. *J Anat Physiol.*. 12:261-287.
- Miall, L.C. and Greenwood, F. (1879b). The anatomy of the Indian elephant. Part II. Muscles of the head and trunk. *J Anat Physiol.* 12:385-400.

- Miall, L.C. and Greenwood, F. (1879c). The anatomy of the Indian elephant. Part III. *J Anat Physiol.* **13**:17-50. Robertson-Bullock, W. (1962). The weight of the African elephant *Loxodonta africana*. *Proc Zool Soc Lond* **138**:133-135.
- Rooney, J.R., II, Sack, W.O. and Habel, R.F. (1967). Guide to the dissection of the horse. Wolfgang O. Sack, Ithaca, New York.
- Roth, V.L. and Shoshani, J. (1988). Dental identification and age determination in *Elephas maximus*. J Zool (Lond) 214(4):567-588.
- Shimizu, Y., Fujita, T. and Kamiya, T. (1960). Anatomy of a female Indian elephant with special reference to its visceral organs. *Acta Anat Nippon* 35(3):261-301.
- Shindo, T. and Mori, M. (1956a). Musculature of the Indian elephant. Part I. Musculature of the forelimb. Okajimas Folia Anat Japonica 28:89-113.
- Shindo, T. and Mori, M. (1956b). Musculature of the Indian elephant. Part II. Musculature of the hindlimb. Okajimas Folia Anat Japonica 28:114-147.
- Shindo, T. and Mori, M. (1956c). Musculature of the Indian elephant. Part III. Musculature of the trunk, neck, and head. *Okajimas Folia Anat Japonica* 29:17-40.
- Shoshani, J., Alder, R., Andrews, K., Baccala, M.J., Barbish, A., Barry, S., Battiata, R., Bedore, M.P., et al (1982). On the dissection of a female Asian elephant (*Elephas maximus maximus* Linnaeus, 1758) and data from other elephants. *Elephant* 2 (1):3-93.
- Shoshani, J. (1996). Skeletal and other basic anatomical features of elephants. In: *The Proboscidea: evolution and palaeoecology of elephants and their relatives*. Shoshani, J. & Tassy, P. eds., Oxford University Press, Oxford. p. 9-20.
- Sikes, S.K. (1971). The Natural History of the African Elephant. American Elsevier Publishing Company, Inc., New York.
- Sisson and Grossman's the anatomy of the domestic animals. Getty, R., ed. (1975). W.B. Saunders Company, Philadelphia.
- Sreekumar, K.P. and Nirmalan, G. (1989). Estimation of body weight in Indian elephants (*Elephas maximus indicus*). Vet Res Commun 13:3-9.

# Researchers Interested in Participating in Necropsy Procedures

The following researchers may be available to participate in necropsies. If you are interested, please contact them as soon as possible after an animal dies or before euthanasia.

Name	Work Number	Home Number	Fax Number
William A. Akerston, Pocatello, Idaho, USA	(208) 236-4151		(208) 236-4600
D. J. Hillman, Baton Rouge, Louisiana, USA	(504) 346-3246	(504) 272-0156	(504) 346-3329
Vaughan A. Langman Shreveport, Louisiana, USA email: 75554,3044@compuserv.com	(318) 797-5244	(318) 798-2750	(318) 797-5156 (318) 227-1141
Susanne J. Miller, Pocatello, Idaho, USA	(208) 526-0938	(208) 529-2453	(208) 236-4600
L. E. Rasmussen, Beaverton, Oregon, USA	(503) 690-1263	(503) 621-1435	(503) 690-1464
Jeheskel (Hezy) Shoshani, Bloomfield Hills, Michigan, USA	(810) 540-3947	(810) 540-3947	(810) 540-3948

### **Research Requests**

1. Dr. Jeheskel (Hezy) Shoshani
Elephant Research Foundation
Department of Biological Sciences
Wayne State University
Detroit, Michigan 48202
Work: (313) 577-1757

e mail: jshosh@sun.science.wayne.edu

Entire fetus or newborn or measurements and photographs of fetus or newborn (Anatomical Studies) Freeze fetus or preserve in 10% formalin. If entire fetus cannot be obtained, photographs and measurements of teeth are particularly needed.

Intact brain (Anatomical Studies) from any age, sex or species of elephant. Preserve in 10% formalin.

Dr. Bets Rasmussen
 Associate Professor of Chemistry
 Oregon Graduate Institute
 20000 NW Walker Rd
 Beaverton, Oregon 97006

e-mail betsr@admin.ogi.edu

phone

home: 503-621-1435 work: 503-690-1263 Fax: 503-690-1464

For any MALE ASIAN (and or AFRICAN) ELEPHANT please call Bets Rasmussen immediately at 503-690-1263 or 503-621-1435. She will come on the next airplane as certain tissues are of extreme and immediate value in her studies.

Vomeronasal organ (Chemical sense studies; electron microscopy). See diagrams A and B. Adults: Only the incisive ducts are requested. These paired openings on the roof of the mouth are 1-2 cm in diameter, several cms posterior to the juncture of the mucosa of the mouth with the lip region. The incisive ducts (ID) are indicated by the white arrows diagram B. You are looking down at the roof of the mouth with the upper teeth toward the bottom of the photograph. The vomeronasal organ proper is located in the region

indicated by the white "BELOW" on diagram B.

Fetal, neonates & young elephants: The whole organ is requested. It is found in the vomer bone, dorsal and posterior to the incisive ducts. It is paired and eventually each half becomes tubular in shape, and is surrounded by a cartilage sheath. In immediately post-natal elephants the incisive ducts are discontinuous with the actual vomeronasal organ and the organ is less tubular. In these young elephants one way to obtain the organ is to work dorsal/posterior from the ducts, looking for shiny white cartilage surrounding receptive tissue, which is hollow in the center.

Fixatives: Preferred fixative is 4% buffered paraformaldehyde or 2% glutaldehyde in a pH 7.2 buffer. Pieces should be VERY SMALL (1mm x 1mm x 1 mm) for EM material. Investigator will ship pre-made fixative upon request. If those two fixatives are not available, fix in buffered 10% formalin and use larger pieces. Save (in formalin) the larger section from which the EM piece was taken. EXTRA VNO TISSUE SHOULD BE FROZEN IN LIQUID NITROGEN IF POSSIBLE IN 1CM X 1CM X 2CM PIECES, FOR BOTH ADULT AND YOUNG ELEPHANTS.

Palatal pits: (Histological and cytological studies) The palatal pits are a dual series of small openings (0-13), asymmetrically and bilaterally located along the approximate demarcation line in the upper head between the hard palate and the trunk. Push aside the upper lip to locate. They are smaller than the incisive openings. Their location is indicated by PP on diagram B.

Trunk tip finger(s): (Chemical sense studies) Both EM fixed (very small pieces & special fixative, see VNO section) and formalin fixed pieces of the dorsal trunk tip finger and corresponding ventral "heel" of Asians and the dorsal and ventral opposing tips of Africans of ALL ages,

phorynx

crunial pinuses

cribriform plate of ethmold

vino duct


include duct

soft paints

Diagram A

especially young and fetal elephants are requested. Especially to be fixed are pieces including the small invaginations.

Diagram B



#### 3. Dr. C. Earle Pope

Audubon Center for Research of Endangered Species

14001 River Road

New Orleans, Louisiana, USA 70131

Work: (504) 398-3161 Home: (504) 734-5381 Fax: (504) 391-7707 email: epope@acres.org

Intact ovaries (To recover oocytes for in-vitro maturation and culture) Remove ASAP; rinse with saline to remove blood and adhering tissue. Wrap in sterile gauze pre-soaked in saline and place in plastic bag or specimen container. Keep at room temperature if sample can be shipped the same day; if longer pack in crushed ice. Ship overnight; will pay shipping.

#### 4. Vaughan A. Langman Ph.D.

LSU in Shreveport Department of Biological Sciences

8515 Youree Drive

Shreveport, Louisiana, USA 71115

Work: (318) 797-5244 Home: (318) 798-2750 Fax: (318) 797-5156

Or

Dr. D. J. Hillman

Louisiana State University

Department of Veterinary Anatomy

School of Veterinary Medicine

Baton Rouge, Louisiana, USA 79803

Work: (504) 346-3246 / 346-3164

Home: (504) 272-0156 Fax: (504) 346-3329

Intact head, neck and ears with cervical vertebrae if possible (To study venous drainage) Contact prior to necropsy. Will pay shipping.

#### 5. Dr. Larry Agenbroad

The Mammoth Site

1800 Hwy 18 By-pass

Hot Springs, South Dakota, USA 57747

Work: (605) 745-6017 Fax: (605) 523-2626

Skeletal remains; Asian or African (Exhibit; comparative collection) Rough flesh if possible and ship frozen or just freeze. Will pay shipping.

6. Dr. William A. Akerston

Curator of Vertebrate Paleontology Idaho Museum of Natural History

Campus Box 8096

Pocatello, Idaho, USA 83209 Work: (208) 236-4151 / 236-3168

Home: (208) 232-5436 Fax: (208) 236-4600

Or

Susanne J. Miller 1450 Antares Drive

Idaho Falls, Idaho, USA 83402

Work: (208) 526-0938 Home: (208) 529-2453 email work: ski@inel.gov email home: olbonz@srv.net

Complete or partial juvenile skeleton (Aid identification of fossil mammoths for comparative osteology collection) Bones must be rough-fleshed and dried to avoid spoilage during shipping. Will pay shipping costs and labor for rough-fleshing.

7. Dr. Linda Munson

Department of Veterinary Pathology, Microbiology Immunology University of California School of Veterinary Medicine

Davis, CA 95616 Work: (916) 752-1385

Sections of uterine endometrium (Characterization of endometrial lesions) Endometrial samples including any polyps, cysts, tumors or other lesions. Samples should include lesion and adjacent normal tissue. Fix in 10% formalin. Ship by U.S. mail. Will pay shipping.

Jenifer Lienau or Carla Bitter
 Arizona Science Center
 600 E. Washington Street

Phoenix, Arizona, USA 85004

Work: (602) 716-2000 Fax: (602) 716-2088

Elephant bladder with ureters and urethra attached. (For museum display) Store in electrolyte solution or 10% saline and freeze in expanded position. Ship frozen, overnight on dry ice. Will pay shipping.

9. Robert L. Van Critter, M.D.

Professor of Medicine

Physiology and Biophysics

University of Washington School of Medicine

Mail: RG-22

University of Washington School of Medicine

Seattle, Washington, USA 98105

Shipping: University of Washington Cardiology

1959 NE Pacific Room E-303 Seattle, Washington, USA 98105

Work: (206) 543-9952 Home: (206) 774-1770 Fax: (206) 543-3639

Cross sections of coronary arteries. (For comparative circulation studies) Take a) 2 pieces, 1 cm in length of right or left main coronary artery; b) 2 pieces of tertiary branch of either coronary artery. Place in formalin and ship by common carrier. Will pay shipping.

10. David Combs, Ph.D.

California Museum of Science and Industry

Director's Office

700 State Drive

Los Angeles, California, USA 90037

Work: (213) 744-7547 Home: (818) 791-5404 Fax: (213) 744-2034

Intact Adult Indian or African elephant heart. (For comparative mammalian blood volume and relative heart sizes museum display) Excised heart must be intact, preferably not opened, leaving 4" or 5" of the aorta and vena cavae as well as 2" of the pulmonary vessels. Pack in ice and ship overnight. Will pay shipping.

11. Drs. Dick Montali and Laura K. Richman

National Zoological Park, Department of Pathology

3001 Block of Connecticut Avenue, NW

Washington, DC, USA 20008-2598

Work: (202) 673-4869

Home: (301) 926-2622 (RJM); (301) 253-8723 (LKR)

Fax: (202) 673-4660

1. Complete set of formalin fixed tissues (as per SSP Necropsy Protocol) 2. Ultra frozen (-70°c) heart, liver, tongue, spleen, intestinal or oral ulcers. 3. Whole blood (20 cc); serum (20 cc). 4. 20 cc urine. (For elephant herpes virus study) For formalin fixed tissues .5 - 1.0 cm thick sections. Take samples to be frozen as soon as possible - place in sterile container and freeze. Send frozen tissue overnight on dry ice. Send blood and urine samples overnight on wet or dry ice. Call before shipping. Will pay shipping.

Dr. Dick Montali / Dr. Jamie Brown
 National Zoological Park, Department of Pathology
 3001 Block of Connecticut Avenue, NW
 Washington, DC, USA 20008-2598

Work: (202) 673-4869 Home: (301) 926-2622 Fax: (202) 673-4660

- 1. Sections of uterine leiomyoma. 2. Sections of normal uterine wall. 3. Serum sample (2-3 ml). (For comparative study of nuclear receptors and estrogen receptors in tumored and normal uterus with human and guinea pig models to identify non GnRH analogs to reverse tumors) Need 2-3 representative sections of leiomyomas and normal uterine wall approximately 1-2 cm thick. Fix 1 set in 10% neutral buffered formalin. Freeze a replicate set in liquid nitrogen or "Revco" ultra freezer at -70°C. Send frozen samples overnight on dry ice. Will pay shipping. Call for billing number. Please include brief reproductive history.
- 13. Anthony T. Boldurian, Ph.D.
  Associate Professor Anthropology
  University of Pittsburgh at Greensburg
  Smith Science Building
  1150 Mr. Pleasant Road
  Greensburg, Pennsylvania, USA 15601

Work: (412) 836-9989 Fax: (412) 836-7129

Two whole large thoracic ribs. (For experimental archeological study to replicate mammoth shaft wrench artifact) Minimum width 60 mm; minimum thickness 20 mm. Samples must be in a "green" or unweathered state, preferably from a recently deceased individual. Pack in ice if tissue still adhering. Call before shipping. Will pay shipping.

14. Dr. Ellen DierenfeldWildlife Conservation Society185 th Street and Southern Blvd.

Bronx, NY 10460 Work: (718) 220-7102 Home: (718) 885-0402 Fax: (718) 220-7126

Email: edierenfeld@wcs.org

Liver, adipose tissue, skeletal muscle, heart. (Baseline data collection on fatty acids, minerals and vitamin concentrations in tissues for comparative purposes.) Take 5 gram samples of each tissue, trim and place in individual plastic bags labeled with the animal's identification, institution, date and tissue. Freeze. Ship on dry ice in hot months, wet ice in cold months in a well-insulated container, overnight. Call for shipping account number.

#### 15. Mark Prausnitz

Gerogin Institute of Technology School of Chemical Engineering Atlanta, GA 30332-0100

Work: (404) 894-5135 Home: (404)92-4597 Fax: (404) 894-2866

Email: mark.prausnitz@che.gatech.edu

Skin. (To study the effects of electric fields on skin from large land mammals as a method to increase delivery of drugs across the skin.) Take full thickness skin samples 12" X 12" from the back, abdomen and chest. Submit 2 - 3 pieces from each site. Skin should have as little hair as possible and not be physically damaged. Animal should be older than 1 year with no known skin diseases. If cadaver is stored refrigerated, skin should be removed within 1 week post mortem. If the cadaver is unrefrigerated, skin should be removed within 1 day. Store the skin frozen, preferably at -60° to -90°C, a conventional freezer is also acceptable. Skin can be stored frozen for up to a week before shipping. Label site and place samples in individual zip-lock bags. Ship in a styrofoam container on dry ice overnight. Will pay shipping.